

On distinguishing special trees by their chromatic symmetric functions

Melanie Gerling

February 28, 2017

Abstract

In 1995, Stanley introduced the well-known chromatic symmetric function $X_G(x_1, x_2, \dots)$ of a graph G . It is a sum of monomial symmetric functions such that for each vertex coloring of G there is exactly one of these summands. The question, whether $X_G(x_1, x_2, \dots)$ distinguishes nonisomorphic trees with the same number of vertices, is still open in general. For special trees it has already been shown. In 2008, Martin, Morin and Wagner proved it for spiders and some caterpillars. We decompose a tree by separating the leafs and their neighbors and do the same to the remaining forest until there remains a forest with vertices of degree not greater than 1. For nonisomorphic trees G and H with the same number of vertices and special properties concerning their number of leafs and leaf neighbors for each subgraph in their leaf decomposition we prove that the chromatic symmetric function distinguishes the graphs. Our idea is to find independent partitions of G and H with respectively a block of maximal cardinality for G as well as for H . These cardinalities are different for graphs G and H with special properties of their leaf decompositions. Additionally, we give explicit formulas for the cardinality of such a maximal block of an independent partition of star connections and spiders.

1 Introduction

Let M be a set and let $\mathcal{P}(M) = \{X \mid X \subseteq M\}$ denote the power set of M . For a natural number $k \geq 1$ let $\mathcal{P}_k(M) = \{X \in \mathcal{P}(M) \mid |X| = k\}$. Because we will exclude certain types of graphs, a simplified definition of a graph suffices. We define a graph G as a tuple $G = (V(G), E(G)) = (V, E)$ of two sets $V(G) = V$ and $E(G) = E$ with $E \subseteq \mathcal{P}_2(V)$. The members of V are called **vertices** and the members of E **edges** of G . This definition excludes the occurrence of **loops**, i. e. members of $E \cap \mathcal{P}_1(V)$, and **multiple edges**, which are irrelevant for our considerations. Two

vertices $v_1, v_2 \in V$ are called **adjacent** to each other, if $\{v_1, v_2\} \in E$. For a vertex $v \in V$ the number of all vertices that are adjacent to v in G is called the **degree** of v in G and denoted by $\deg_G(v)$.

A graph $H = (V', E')$ is called a **subgraph** of $G = (V, E)$, if $V' \subseteq V$ and $E' \subseteq E$. We also write $H \subseteq G$. A subgraph is an **induced subgraph** of G , if $E' = E \cap \mathcal{P}_2(V')$. For $X \subseteq V$ the graph $G[X]$ is the subgraph induced by the set X and for $\{v_1, \dots, v_r\} \subseteq V$ we set $G - \{v_1, \dots, v_r\} = G[V \setminus \{v_1, \dots, v_r\}]$.

Let $G = (V, E)$ be a finite graph and $n = |V|$. A function $\kappa : V \rightarrow Y$ with a set of colors $Y = \{1, 2, \dots\}$ is called a **vertex coloring**. It is called a **proper vertex coloring**, if for all edges $\{v_1, v_2\}$ the relation $\kappa(v_1) \neq \kappa(v_2)$ holds. The number of all proper vertex colorings for a graph G in at most y colors was introduced as a polynomial $P(G; y)$ by [1] and was called the **chromatic polynomial** by [2]. Now, we turn to a similar function.

Definition 1 ([5], p. 168)

Let $x_1, x_2, \dots$ be (commuting) indeterminates and $V = \{v_1, v_2, \dots, v_n\}$. Then the **symmetric function generalization of the chromatic polynomial** of G is defined as

$$X_G(x_1, x_2, \dots) = \sum_{\kappa} x_{\kappa(v_1)} x_{\kappa(v_2)} \dots x_{\kappa(v_n)}$$

where $\kappa : V \rightarrow Y$ is a proper vertex coloring.

This function has the following connection with the chromatic polynomial.

Remark 2 ([5], p. 169)

For $r \in Y$ we have $X_G(1^r) = P(G; r)$.

For the next Remark we need some definitions.

Definition 3 Let $G = (V, E)$ be a graph. We call a partition of V a **partition of G** . $\Pi(G)$ denotes the set of all partitions of G . Each member of a partition is called a **block**.

We will only be interested in partitions which are defined as follows.

Definition 4 Non adjacent vertices and edges which don't have any vertex in common are called **independent of each other**. A set $X \subseteq V$ or $Y \subseteq E$ is said to be **independent** if its members are pairwise independent of each other.

An **independent partition** π of $G = (V, E)$ denotes, as defined in [3], section 3, a partition of G , such that each block $A \in \pi$ is independent. We denote the set of all independent partitions by $\Pi_I(G)$.

The next observation will be very useful to answer the question for special types of trees, whether two not isomorphic trees with the same number of vertices can be distinguished by their symmetric function generalization of the chromatic polynomial or not.

Remark 5 We can identify a vertex coloring of a graph $G = (V, E)$ with an independent partition of V , such that the vertices of each block are colored in the same color and vertices from different blocks are colored in different colors. As Definition 1 shows, each summand $x_{\kappa(v_1)}x_{\kappa(v_2)} \dots x_{\kappa(v_n)}$ contains a product $x_{\kappa(v_i)}^k$, if there are vertices $v_1, \dots, v_k \in V$ with $\kappa(v_1) = \dots = \kappa(v_k)$, which means that they are in the same block. So, k shows the cardinality of that block.

For special types of non-isomorphic trees with the same number of vertices it will be easy to show, that the cardinalities of their blocks with maximal cardinality are different from each other. So the chromatic symmetric functions of these trees differ of course, too.

We call a graph $P = (V, E)$ with vertex set $V = \{v_0, \dots, v_k\}$ and edge set $E = \{\{v_0, v_1\}, \{v_1, v_2\}, \dots, \{v_{k-1}, v_k\}\}$ with $|V| \geq 1$ and pairwise distinct v_i for $i = 0, \dots, k$ a **path** of length k from v_0 to v_k .

A non-empty graph is called **connected**, if for all $u, w \in V$ there is a path from u to w , i. e. a path P with $u = v_0$ and $w = v_k$.

A graph F without any cycles is called a **forest**, a connected forest T is called a **tree** and a vertex $v \in V(F)$ of degree one is called a **leaf**.

Definition 6 Let $T = T_1 = F_1$ be a tree, the vertices $v_1, v_2, \dots, v_{b_1} \in V(T)$ its leaves and $w_1, w_2, \dots, w_{\eta_1} \in V(T)$ all vertices that are adjacent to a leaf. Of course, $b_1 \geq \eta_1$ and $F_1 - \{v_1, \dots, v_{b_1}, w_1, \dots, w_{\eta_1}\} = F_2$ is a forest. For F_2 we do the same and continue in this manner. We get a subgraph relation $T_1 = F_1 \subseteq F_2 \subseteq \dots \subseteq F_k$, such that F_k is a forest with vertex degrees ≤ 1 . For each $i \in \{1, \dots, k-1\}$ there are leaves $v_{i1}, v_{i2}, \dots, v_{i_{b_i-1}} \in V(F_i)$ and all possible vertices $w_{i1}, w_{i2}, \dots, w_{i_{\eta_i-1}} \in V(F_i)$, that are adjacent to a leaf. If in the forest F_k all vertices have degree 0, we have $b_1 \geq \eta_1 \geq b_2 \geq \eta_2 \geq \dots \geq b_k \geq \eta_k$ with $\eta_k = 0$. But if there are α vertices $v_{kb_k1}, \dots, v_{kb_k\alpha}$ of degree 1 with $1 \leq \alpha \leq |V(F_k)|$, we set $b_k = |V(F_k)| - \frac{\alpha}{2}$ and $\eta_k = \frac{\alpha}{2}$. Note, that α is always even. Then $b_1 \geq \eta_1 \geq b_2 \geq \eta_2 \geq \dots \geq b_k \geq \eta_k$ holds. For a tree T we call such a relation a **leaf decomposition** and denote it by (T, k) .

Remark 7 For every tree T there is exactly one $k \in \mathbb{N}$, $k \geq 0$, such that (T, k) is a leaf decomposition of T as in Definition 6.

Each vertex $v \in V(T)$ is of the form v_{ij_1} or w_{ij_2} with $i \in \{1, \dots, k\}$, $j_1 \in \{1, \dots, b_i\}$ and $j_2 \in \{1, \dots, \eta_i\}$. So, v is either a leaf or a neighbor of a leaf in the associated graph F_i .

Note, that two non-isomorphic trees are not determined by their tree decomposition as the following example shows.

Example 8 In the following picture there are two non-isomorphic trees T_1 and T_2 with the same leaf decomposition. For T_1 we define the leaf decomposition $b_{11} \geq \eta_{11} \geq b_{12}$ and for T_2 the leaf decomposition $b_{21} \geq \eta_{21} \geq b_{22}$. We have $b_{11} = b_{21} = 6$, $\eta_{11} = \eta_{21} = 3$ and $b_{12} = b_{22} = 1$.




Figure 1: T_1 and T_2

2 The number of leaves and further properties

Theorem 9 Let $n \geq 3$. For a tree T with $|V(T)| = n$ there is a maximum $|B|$ with $B \in \sigma$, $\sigma \in \Pi_I(T)$, such that for all leaves $v \in V(T)$ we have $v \in B$.

Proof:

For all leaves $v \in V(T)$ there is a vertex w with $N_T(v) = \{w\}$, i. e. of course $|N_T(v)| = 1$. From all such pairs $\{v, w\}$ we may choose at most one vertex for our block B because of its independence and because of the maximality of $|B|$ we have

to chose at least one. We choose always the leaf v because the number of neighbors of leafs is equal to or smaller then the number of leafs. So we have $v \in B$ for all leaves $v \in V(T)$.
q.e.d.

The number of leaves can be an important hint to decide, whether two trees can be distinguished by their symmetric function generalization of the chromatic polynomial as the next theorem shows.

Theorem 10 *Let T_1 and T_2 be trees with $|V(T_1)| = |V(T_2)| = n \geq 4$. The number of leaves of T_i for $i \in \{1, 2\}$ is denoted by b_i and $b_1 > b_2$. Let η_i be the number of all vertices w with $v \in N_{T_i}(w)$ for a leaf $v \in V(T_i)$ and let $\rho_i = n - \eta_i - b_i$. Let $V(\rho_i)$ the set of vertices which contribute to ρ_i and let $T[V(\rho_i)]$ be a path. Furthermore, we assume one of the following cases:*

1. $\rho_1 = \rho_2$,
2. $\rho_1 > \rho_2$,
3. $\rho_1 < \rho_2$ and $b_1 - b_2 > \lceil \frac{\rho_2 - \rho_1}{2} \rceil$,
4. $\rho_1 < \rho_2$ and $b_1 - b_2 > \lceil \frac{\rho_1 - \rho_2}{k} \rceil$ for all $k \in \mathbb{N}$ with $k \neq 2$ and $\frac{k}{k-2} \leq \rho_2 - \rho_1 < k(b_1 - b_2)$.

Then $X_{T_1}(x_1, x_2, \dots) \neq X_{T_2}(x_1, x_2, \dots)$.

Proof:

Because of Theorem 9 we compare independent partitions of T_1 and T_2 , such that there is respectively one block B_i with $|B_i|$ maximum and $v_i \in B_i$ for all leaves $v_i \in V(T_i)$. We set

$$M_i = \max\{|B_i| \mid B_i \in \sigma_i \wedge \sigma_i \in \Pi_I(T_i)\}.$$

For each case we will show $M_1 > M_2$. Note, that we can choose from the vertices ρ_i exactly $\lceil \frac{\rho_i}{2} \rceil$ further vertices for the block B_i because we did not choose any neighbor of a leaf and because of the path structure of $T[V(\rho_i)]$.

1. (a) Case ρ_1 is even. Because of $b_1 > b_2$ and $\rho_1 = \rho_2$ we have

$$M_1 = b_1 + \frac{\rho_1}{2} > b_2 + \frac{\rho_2}{2} = M_2.$$

- (b) Case ρ_1 is odd. Like in the first case we get

$$M_1 = b_1 + \frac{\rho_1 + 1}{2} > b_2 + \frac{\rho_2 + 1}{2} = M_2.$$

2. (a) Case ρ_1 and ρ_2 are even. Because of $b_1 > b_2$ and $\rho_1 > \rho_2$ it leads to

$$M_1 = b_1 + \frac{\rho_1}{2} > b_2 + \frac{\rho_2}{2} = M_2 .$$

- (b) Case ρ_1 and ρ_2 are odd. Like above the following holds:

$$M_1 = b_1 + \frac{\rho_1 + 1}{2} > b_2 + \frac{\rho_2 + 1}{2} = M_2 .$$

- (c) Case ρ_1 is odd and ρ_2 is even. Similar as above we get

$$M_1 = b_1 + \frac{\rho_1 + 1}{2} > b_2 + \frac{\rho_2}{2} = M_2 .$$

- (d) Case ρ_1 is even and ρ_2 is odd. Like above and because of $\rho_1 \geq \rho_2 + 1$ it leads to

$$M_1 = b_1 + \frac{\rho_1}{2} > b_2 + \frac{\rho_2 + 1}{2} = M_2 .$$

3. (a) Case ρ_1 and ρ_2 are even. Hence,

$$b_1 - b_2 > \left\lceil \frac{\rho_2 - \rho_1}{2} \right\rceil$$

$$M_1 = b_1 + \frac{\rho_1}{2} > b_2 + \frac{\rho_2}{2} = M_2 .$$

- (b) Case ρ_1 and ρ_2 are odd. Similary we get

$$b_1 - b_2 > \left\lceil \frac{\rho_2 - \rho_1}{2} \right\rceil$$

$$M_1 = b_1 + \frac{\rho_1 + 1}{2} > b_2 + \frac{\rho_2 + 1}{2} = M_2 .$$

- (c) Case ρ_1 is odd and ρ_2 is even. Like above we have

$$b_1 - b_2 > \frac{\rho_2 - \rho_1}{2} - \frac{1}{2}$$

$$M_1 = b_1 + \frac{\rho_1 + 1}{2} > b_2 + \frac{\rho_2}{2} = M_2 .$$

- (d) Case ρ_1 is even and ρ_2 is odd. Thus,

$$b_1 - b_2 > \left\lceil \frac{\rho_2 - \rho_1}{2} \right\rceil$$

$$b_1 - b_2 > \frac{\rho_2 - \rho_1}{2} + \frac{1}{2}$$

$$M_1 = b_1 + \frac{\rho_1}{2} > b_2 + \frac{\rho_2 + 1}{2} = M_2 .$$

4. First we show that $\frac{\rho_2 - \rho_1}{k} \geq \frac{\rho_2 - \rho_1}{2} + \frac{1}{2}$ holds. As assumed, we have

$$\begin{aligned}
\rho_2 - \rho_1 &\geq \frac{k}{k-2} \\
\Leftrightarrow (k-2)(\rho_2 - \rho_1) - k &\geq 0 \\
\Leftrightarrow (2-k)\rho_2 + (k-2)\rho_1 - k &\geq 0 \\
\Leftrightarrow 2\rho_2 - 2\rho_1 &\geq k\rho_2 - k\rho_1 + k \\
\Leftrightarrow \frac{\rho_2 - \rho_1}{k} &\geq \frac{\rho_2 - \rho_1}{2} + \frac{1}{2}.
\end{aligned}$$

(a) Case ρ_1 and ρ_2 are even. Then we get

$$\begin{aligned}
b_1 - b_2 &\geq \left\lceil \frac{\rho_2 - \rho_1}{k} \right\rceil \geq \left\lceil \frac{\rho_2 - \rho_1}{2} \right\rceil \\
M_1 = b_1 + \frac{\rho_1}{2} &\geq b_2 + \frac{\rho_2}{2} = M_2.
\end{aligned}$$

(b) Case ρ_1 and ρ_2 are odd. It leads to

$$\begin{aligned}
b_1 - b_2 &\geq \left\lceil \frac{\rho_2 - \rho_1}{k} \right\rceil \geq \left\lceil \frac{\rho_2 - \rho_1}{2} \right\rceil \\
b_1 - b_2 &\geq \frac{\rho_2 + 1}{2} - \frac{\rho_1 + 1}{2} \\
M_1 = b_1 + \frac{\rho_1 + 1}{2} &\geq b_2 + \frac{\rho_2 + 1}{2} = M_2.
\end{aligned}$$

(c) Case ρ_1 is odd and ρ_2 is even. Similary as above we have

$$\begin{aligned}
b_1 - b_2 &\geq \left\lceil \frac{\rho_2 - \rho_1}{k} \right\rceil \geq \left\lceil \frac{\rho_2 - \rho_1}{2} \right\rceil \\
b_1 - b_2 &\geq \frac{\rho_2 - \rho_1}{2} - \frac{1}{2} \\
M_1 = b_1 + \frac{\rho_1 + 1}{2} &\geq b_2 + \frac{\rho_2}{2} = M_2.
\end{aligned}$$

(d) Case ρ_1 is even and ρ_2 is odd. Hence,

$$\begin{aligned}
b_1 - b_2 &\geq \left\lceil \frac{\rho_2 - \rho_1}{k} \right\rceil \geq \left\lceil \frac{\rho_2 - \rho_1}{2} \right\rceil \\
b_1 - b_2 &\geq \frac{\rho_2 - \rho_1}{2} + \frac{1}{2} \\
M_1 = b_1 + \frac{\rho_1}{2} &\geq b_2 + \frac{\rho_2 + 1}{2} = M_2.
\end{aligned}$$

q.e.d.

We show an example for Theorem 10. It contains two graphs which are special cases of the graphs claimed in Theorem 10. This special graph type was already introduced by [4]. They already proved that so-called caterpillars can be distinguished by their chromatic symmetric functions.

Example 11 *Let $n = 4$. We consider the following two graphs. On the left side there is the graph S_4 and on the right side we see the graph P_4 . We have $|V(S_4)| = |V(P_4)| = 4$. For the graph S_4 we have $b_1 = 3$, $\eta_1 = 1$ and $\rho_1 = 0$, while we have*




Figure 2: S_4 and P_4

$b_2 = 2$, $\eta_2 = 2$ and $\rho_2 = 0$ for the graph P_4 . Because of $b_1 > b_2$ and $\rho_1 = \rho_2$ Theorem 10 leads to $3 = M_1 > M_2 = 2$ as we can see easily.

Now, we will pay attention to a special situation which is interesting for our further considerations.

Situation 12 *Let T_{11} and T_{21} be trees with $|V(T_{11})| = |V(T_{21})| = n$ and the leaf decompositions (T_{11}, k_1) and (T_{21}, k_2) . We set $r = \max \{k_1, k_2\}$. Let $T_{11} \subseteq F_{12} \subseteq F_{13} \subseteq \dots \subseteq F_{1k_1}$, and for $i \in \{1, \dots, k_1\}$ there exist respectively b_{1i} leafs and η_{1i} neighbor vertices like in Definition 6. Analogously, let $T_{21} \subseteq F_{22} \subseteq F_{23} \subseteq \dots \subseteq F_{2k_2}$ with b_{2j} and η_{2j} for $j \in \{1, \dots, k_2\}$ like in Definition 6.*

We set $(T_{11}, k_1) = (T_{11}, r)$ and $(T_{21}, k_2) = (T_{21}, r)$, where b_{1i} and η_{1i} or b_{2j} and η_{2j} become zero for $k_1 \neq k_2$ from a certain i or respectively j on.

We turn to some special cases. The next theorem shows the first one.

Theorem 13 *We regard Situation 12 for trees T_1 and T_2 and assume $\eta_{1i} \leq \eta_{2i}$ as well as $b_{1i} \geq b_{2i}$ for all $i \in \{1, \dots, r\}$. We except the case that there is equality everywhere. Then $X_{T_1}(x_1, \dots) \neq X_{T_2}(x_1, \dots)$.*

Proof:

By Theorem 9, for $k = 1, 2$ there is a partition $\sigma_k \in \Pi_I(T_k)$ with $B_k \in \sigma_k$ and $|B_k|$ maximal, such that for all leaves $v \in T_k$ we have $v \in B_k$. Let M_k as in the proof of Theorem 10. Then we have $b_{ki} \in B_k$, $\eta_{ki} \notin B_k$ for all $i \in \{1, \dots, r\}$. Because we have at least one strict inequality the relation $M_1 > M_2$ holds because of $|V(T_1)| = |V(T_2)| = n$ and so $X_{T_1}(x_1, \dots) \neq X_{T_2}(x_1, \dots)$.
q.e.d.

Example 14 *Let $n = 8$. We consider the following two graphs. On the left side there is a tree T_1 and on the right side we see another tree T_2 . We have $|V(T_1)| = |V(T_2)| = 8$. We get the following relations:*




Figure 3: T_1 and T_2

$$\begin{aligned}
b_{11} = 5 &> b_{21} = 4 \\
\eta_{11} = 2 &< \eta_{21} = 4 \\
b_{12} = 1 &> b_{22} = 0 \\
\eta_{12} = 0 &= \eta_{22} = 0,
\end{aligned}$$

such that Theorem 13 holds. Indeed,

$$|M_1| = 5 + 1 = 6 > |M_2| = 4 + 0 = 4.$$

In the following Theorem we introduce another case.

Theorem 15 *We consider again Situation 12. Let $i_1, i_2, \dots, i_s \in \{1, 2, \dots, r\}$, $1 \leq s \leq r-1$ and $b_{1j} > b_{2j}$ for all $j \in \{1, 2, \dots, r\} \setminus \{i_1, i_2, \dots, i_s\}$ and $b_{1i_t} \leq b_{2i_t}$ for all $t \in \{1, 2, \dots, s\}$. Furthermore,*

$$\sum_{\alpha=i_1}^{i_s} (b_{2\alpha} - b_{1\alpha}) < \sum_{j \in \{1, \dots, r\} \setminus \{i_1, \dots, i_s\}} (b_{1j} - b_{2j}) \quad (1)$$

holds. Then we have $X_{T_1}(x_1, \dots) \neq X_{T_2}(x_1, \dots)$.

Proof:

As in the proof of Theorem 13, we look at $\sigma_k \in \Pi_I(T_k)$ for $k = 1, 2$ with $B_k \in \sigma_k$ and $|B_k|$ maximal, such that for all leaves $v \in T_k$ we have $v \in B_k$. Again, $b_{ki} \in B_k$, $\eta_{ki} \notin B_k$ for all $i \in \{1, \dots, r\}$. Because of inequality 1 we have $M_1 > M_2$ and hence $X_{T_1}(x_1, \dots) \neq X_{T_2}(x_1, \dots)$.
q.e.d.

3 Maximal blocks of independent partitions of star connections

In this section we will be concerned with maximal blocks of independent partitions of a special type of graph. It is constructed from stars, which are renownedly defined as follows.

Definition 16 *A **star** S_n for a natural number $n \geq 2$ is a graph with vertex set $V = \{v_1, \dots, v_n\}$ and edge set $E = \{\{v_1, v_2\}, \{v_1, v_3\}, \dots, \{v_1, v_n\}\}$. The vertex v_1 is called the **center** of the star.*

Now, we introduce two operations. The first one is a graph operation and the second one a set operation.

Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two graphs. Then we can create a new graph $G_1 \cup G_2 = (V_1 \cup V_2, E_1 \cup E_2)$. Note, that the sets V_1 and V_2 (and thus of course possibly E_1 and E_2) are not necessarily disjoint.

An intersection $\bigcap_{i \in \{1, \dots, s\}} A_i$ of sets, such that the $i \in \{1, \dots, s\}$ are pairwise distinct from each other, will be denoted by $\bigcap_{i \in \{1, \dots, s\}}^\bullet A_i$. For a special type of graph we give the following Definition.

Definition 17 Let $S_{n_1}, \dots, S_{n_r}$ be stars with $|V(S_{n_i})| = n_i$ vertices for $i = 1, \dots, r$ with $r \geq 2$ and $n_i \geq 3$. We define a tree $T = (V, E) = \bigcup_{i=1}^r S_{n_i}$ with the following properties: Let C be the set of all centers of stars in T and

$$\begin{aligned} \{v_1, \dots, v_t\} = \\ \left\{ v \in V \setminus C \mid \exists S_{n_{i_1}}, \dots, S_{n_{i_s}} \text{ with } i_1, \dots, i_s \in \{1, \dots, r\}, \{v\} = \bigcap_{l \in \{i_1, \dots, i_s\}}^\bullet S_{n_l} \right\}. \end{aligned}$$

Then we assume $|S_{n_i} \cap S_{n_j}| \leq 1$ for all pairwise distinct $i, j \in \{1, \dots, r\}$ and that no center belongs to more than one star as a center or another vertex. Hence, for each $v \in V \setminus (C \cup \{v_1, \dots, v_t\})$ we have $\deg_G(v) = 1$ and for $v \in \{v_1, \dots, v_t\}$ with $\{v\} = \bigcap_{l \in \{i_1, \dots, i_s\}}^\bullet S_{n_l}$ we have $\deg_G(v) = s$. Then we call the graph T a **star connection** and denote it by $(S_{n_1}, \dots, S_{n_r}, v_1, \dots, v_t)$.

Before we continue with the symmetric function generalization of the chromatic polynomial, we observe the following property of a star connection.

Theorem 18 Like in the proof of Theorem 10, we define

$$M = \max \{ |B_j| \mid B_j \in \sigma_j \wedge \sigma_j \in \Pi_I(T) \}.$$

In the situation of Definition 17 we have

$$M = \sum_{k=1}^r (n_k - 1) - \sum_{i=1}^t (\deg_T(v_i) - 1).$$

Proof:

We will prove the Theorem by complete induction.

For $r = 2$ the graph T is a star connection $S_{n_1} \cup S_{n_2}$ with exactly one common vertex v_1 , that is not a center of a star. Thus, $\deg_G(v_1) = 2$. The greatest set of pairwise

independent vertices is of course the set A of all vertices without the two centers. Additionally, we may count v_1 only one time. Hence, for $|A| = M$ we have

$$\begin{aligned} M &= n_1 + n_2 - 3 \\ &= n_1 + n_2 - 2 - (2 - 1) \\ &= \sum_{k=1}^2 (n_k - 1) - \sum_{i=1}^1 (deg_T(v_i) - 1). \end{aligned}$$

For the induction step we assume that the claim is true for an $r \geq 2$. Let $T = \bigcup_{i=1}^r S_{n_i}$. We want to connect T with a star $S_{n_{r+1}}$ to a new star connection H . Then there is a leaf $w_1 \in V(S_{n_{r+1}})$ which is also a leaf of stars $S_{n_{i_1}}, \dots, S_{n_{i_s}}$ in T with $i_1, \dots, i_s \in \{1, \dots, r\}$ and $1 \leq s \leq r$. We consider the maximal set A_T of pairwise independent vertices in the graph T . Because w_1 is not a center we have $w_1 \in A_T$. By induction, for $|A_T| = M_T$ we get

$$M_T = \sum_{k=1}^r (n_k - 1) - \sum_{i=1}^t (deg_T(v_i) - 1).$$

All the other leaves $w_2, \dots, w_{n_{r+1}}$ of $S_{n_{r+1}}$ are not leaves of other stars. Hence, there are $n_{r+1} - 2$ new vertices in H that are contained in the maximal set A_H of pairwise independent vertices in H . We have $deg_H(w_1) = deg_T(w_1) + 1$.

Case 1: w_1 is identified in H with a leaf of T that is not a connection vertex in T . We set $w_1 = v_{t+1}$. Then $deg_T(v_i) = deg_H(v_i)$ for all $i \in \{1, \dots, t\}$ while $deg_T(v_{t+1}) = 1$ and $deg_H(v_{t+1}) = 2$, such that

$$\begin{aligned} M_H &= M_T + (n_{r+1} - 2) \\ &= \sum_{k=1}^r (n_k - 1) - \sum_{i=1}^t (deg_T(v_i) - 1) + (n_{r+1} - 2) \\ &= \sum_{k=1}^{r+1} (n_k - 1) - \sum_{i=1}^t (deg_T(v_i) - 1) - 1 \\ &= \sum_{k=1}^{r+1} (n_k - 1) - \sum_{i=1}^t (deg_H(v_i) - 1) - (deg_H(v_{t+1}) - 1) \\ &= \sum_{k=1}^{r+1} (n_k - 1) - \sum_{i=1}^{t+1} (deg_H(v_i) - 1). \end{aligned}$$

Case 2: w_1 is identified in H with a vertex that is not a center and not a leaf of T . Then that vertex is already a connection vertex in T , say v_t . As in case 1, $deg_T(v_i) = deg_H(v_i)$ for all $i \in \{1, \dots, t-1\}$ and $deg_T(v_t) + 1 = deg_H(v_t)$, such

that

$$\begin{aligned}
M_H &= M_T + (n_{r+1} - 2) \\
&= \sum_{k=1}^r (n_k - 1) - \sum_{i=1}^t (\deg_T(v_i) - 1) + (n_{r+1} - 2) \\
&= \sum_{k=1}^{r+1} (n_k - 1) - \sum_{i=1}^t (\deg_T(v_i) - 1) - 1 \\
&= \sum_{k=1}^{r+1} (n_k - 1) - \sum_{i=1}^t (\deg_H(v_i) - 1) .
\end{aligned}$$

q.e.d.

Now we will give an example for Theorem 18.

Example 19 We look at the graph $T = S_4 \cup S_5 \cup S_3 \cup S_4$ with $|V(T)| = 13$ vertices, the number of connection vertices $t = 3$ and the number of centers $C = 4$ which is depicted below. In this graph we have $r = 4$, and for the connection vertices




Figure 4: T

$v_1, \dots, v_3$ we can see that $\deg_T(v_1) = \deg_T(v_2) = \deg_T(v_3) = 2$. This leads to

$$\begin{aligned} M &= \sum_{k=1}^r (n_k - 1) - \sum_{i=1}^t (\deg_T(v_i) - 1) \\ &= (3 + 4 + 2 + 3) - (1 + 1 + 1) \\ &= 9. \end{aligned}$$

Now we can easily compare the symmetric function generalizations of the chromatic polynomial of two non-isomorphic star connections with the same number of vertices and get the following result.

Theorem 20 *Let $T_1 = (S_{n_1}, \dots, S_{n_r}, v_1, \dots, v_t)$ and $T_2 = (S_{m_1}, \dots, S_{m_s}, w_1, \dots, w_u)$ two star connections with $|V(T_1)| = |V(T_2)|$ and $r < s$. Then $X_{T_1}(x_1, \dots) \neq X_{T_2}(x_1, \dots)$.*

To prove Theorem 20 we need a Lemma.

Lemma 21 *In the situation of Definition 17 the two following relations hold.*

- a) $|V(T)| = (\sum_{k=1}^r n_k) - (r - 1)$.
- b) $\sum_{i=1}^t (\deg_T(v_i) - 1) = r - 1$.

Proof:

- a) Because of $V(T) = V(\bigcup_{i=1}^r S_{n_i})$ we have just to sum up all n_i for all $i \in \{1, \dots, r\}$ and to subtract all $r - 1$ multiple countings of vertices that connect different stars.
- b) We show $\sum_{i=1}^t \deg_T(v_i) = t + r - 1$.
For each vertex v_i with $i \in \{1, \dots, t\}$ we have already the contribution $1 = \deg_{S_{n_i}}(v_i)$, altogether t . If we construct T step by step by the series $S_{n_1}, S_{n_1} \cup S_{n_2}, \dots, S_{n_1} \cup \dots \cup S_{n_r} = T$, we have $r - 1$ steps. Each of them increases the degree of one of the vertices $v_1, \dots, v_t$ by 1. Thus, $\sum_{i=1}^t \deg_T(v_i) = t + r - 1$.

q.e.d.

Now, we are able to prove Theorem 20.

Proof of Theorem 20:

We show $M_1 > M_2$.

By Theorem 18 and Lemma 21 (b) we have

$$\begin{aligned} M_1 &= \sum_{k=1}^r (n_k - 1) - \sum_{i=1}^t (\deg_{T_1}(v_i) - 1) \\ &= \sum_{k=1}^r (n_k - 1) - (r - 1) \\ &= \left(\sum_{k=1}^r n_k \right) - 2r + 1 \end{aligned}$$

and

$$\begin{aligned} M_2 &= \sum_{k=1}^s (m_k - 1) - \sum_{i=1}^u (\deg_{T_2}(w_i) - 1) \\ &= \sum_{k=1}^s (m_k - 1) - (s - 1) \\ &= \left(\sum_{k=1}^s m_k \right) - 2s + 1 . \end{aligned}$$

By assumption and Lemma 21 (a) we have furthermore

$$|V(T_1)| = \left(\sum_{k=1}^r n_k \right) - (r - 1) = |V(T_2)| = \left(\sum_{k=1}^s m_k \right) - (s - 1) .$$

Because of $r < s$

$$M_1 = \left(\sum_{k=1}^r n_k \right) - 2r + 1 > \left(\sum_{k=1}^s m_k \right) - 2s + 1 = M_2$$

follows.

q.e.d.

4 Maximal blocks of independent partitions of spiders

In this section we will turn our attention to so-called spiders which were introduced by [4]. A **spider** is a tree which has exactly one vertex with degree greater than 2.

Situation 22 Let $S = (V, E)$ be a spider and v the vertex with $r = \deg_S(v) > 2$. We denote the set of leafs of S by $\{l_1, \dots, l_r\} \subseteq V$ and the length of the path from l_i to v by L_i for each $i \in \{1, \dots, r\}$. We distinguish between even and odd lengths. Let $s \in \{0, \dots, r\}$ be the number of even lengths and $t \in \{0, \dots, r\}$ be the number of odd lengths such that $r = s + t$. Either one of the two sets of lengths is empty (i. e. for $s = 0$ or $t = 0$) or we have a set $\{L_{11}, \dots, L_{1s}\}$ of even lengths as well as a set $\{L_{21}, \dots, L_{2t}\}$ of odd lengths.

Now, we can easily give an explicit formula for the cardinality of the biggest block of an independent partition of a spider.

Theorem 23 Given the conditions of Situation 22.

i) If all lengths are even, then

$$M = \sum_{k=1}^r \frac{L_k}{2}$$

holds.

ii) If all lengths are odd, then we have

$$M = \sum_{k=1}^r \frac{L_k - 1}{2} + 1.$$

iii) If there are even and odd lengths, then

$$M = \sum_{k=1}^s \frac{L_{1k}}{2} + \sum_{k=1}^t \frac{L_{2k} - 1}{2}$$

holds.

Proof:

i) For all $k \in \{1, \dots, r\}$ let P_k be the path from l_k to v . By assumption, 2 is a divisor of L_k . For $B \in \sigma$, $\sigma \in \Pi_I(S)$ with $|B|$ maximal we start to select vertices in P_k . Let $V(P_k) = \{l_k, v_{k1}, v_{k2}, \dots, v_{kL_k-2}, v\}$. According to Theorem 9, we start with l_k because it is a leaf in S . Then $l_k, v_{k2}, v_{k4}, \dots, v_{kL_k-2} \in B$ and of course $|\{l_k, v_{k2}, v_{k4}, \dots, v_{kL_k-2}\}| = \frac{L_k}{2}$. Because we started with l_k we have $v \notin B$ and hence we do not need to take account of v .

- ii) Here we apply the same method as above. Now, $L_k - 1$ is even and we get the sequence $l_k, v_{k2}, v_{k4}, \dots, v_{kL_k-3} \in B$ with $|\{l_k, v_{k2}, v_{k4}, \dots, v_{kL_k-3}\}| = \frac{L_k-1}{2}$. But now $v \notin N(v_{kL_k-3})$ and thus we also choose $v \in B$. Because $v = \bigcap_{k=1}^r S_k$ we add exactly 1.
- iii) We just combine the first two parts. Because there are even lengths we can not add 1 for v .

q.e.d.

References

- [1] Birkhoff, G. D.: A determinant formula for the number of ways of coloring an map, *Ann. of Math. (2)*, 14(1), 1912, p. 42-46. <http://www.jstor.org/stable/1967597>
- [2] Birkhoff, G. D. and Lewis, D. C.: Chromatic polynomials, *Trans. Amer. Math. Soc.*, 1946, p. 355-451. <http://www.ams.org/journals/tran/1946-060-00/S0002-9947-1946-0018401-4/home.html>
- [3] Dohmen K., Pönitz A. and Tittmann P.: A new two-variable generalization of the chromatic polynomial, *Discrete Math. Theor. Comput. Sci.*, 6, 2003, p. 69-90. <http://www.emis.de/journals/DMTCS/volumes/abstracts/dm060106.abs.html>
- [4] Martin, J. L., Morin, M. and Wagner, J. D.: On distinguishing trees by their chromatic symmetric functions, *Combin. Theory Ser. A*, 115(2), 2008, p. 237-253. <http://www.sciencedirect.com/science/article/pii/S0097316507000775>
- [5] Stanley, R.: A symmetric function generalization of the chromatic polynomial of a graph, *Adv. Math. Ill*, 1995, p. 166-194. <http://www.sciencedirect.com/science/article/pii/S0001870885710201>